



A service of the National Library of Medicine  
and the National Institutes of Health

My NCBI  
[\[Sign In\]](#) [\[Register\]](#)

[All Databases](#)[PubMed](#)[Nucleotide](#)[Protein](#)[Genome](#)[Structure](#)[OMIM](#)[PMC](#)[Journals](#)[Books](#)

Search  for

☒ Limits☐ Preview/Index☐ History☐ Clipboard☐ Details

Limits: **Publication Date to 2000**

Display  Show  Sort by  Send to

All: 1 Review: 0 ☐

About Entrez  
NCBI Toolbar

Text Version

Entrez PubMed  
Overview  
Help | FAQ  
Tutorials  
New/Noteworthy  
E-Utilities

PubMed Services  
Journals Database  
MeSH Database  
Single Citation Matcher  
Batch Citation Matcher  
Clinical Queries  
Special Queries  
LinkOut  
My NCBI

Related Resources  
Order Documents  
NLM Mobile  
NLM Catalog  
NLM Gateway  
TOXNET  
Consumer Health  
Clinical Alerts  
ClinicalTrials.gov  
PubMed Central

☐ 1: [Biochim Biophys Acta](#). 2000 Nov 15;1502(3):337-50. [Related Articles, Links](#)

**ELSEVIER**  
**FULL-TEXT ARTICLE**

## Ovarian tumor cells express a novel multi-domain cell surface serine protease.

Underwood LJ, Shigemasa K, Tanimoto H, Beard JB, Schneider EN, Wang Y, Parmley TH, O'Brien TJ.

Department of Biochemistry and Molecular Biology, University of Arkansas for Medical Sciences, Little Rock, AR 72205-7199, USA.

Serine proteases serve many functions in normal biological processes. These functions are often usurped by cancer cells to allow progression of tumors by increasing the growth and metastatic potential of the neoplasia. Here, we have used a polymerase chain reaction (PCR)-based strategy to clone Tumor Associated Differentially-expressed Gene-12 (TADG-12), a new serine protease from ovarian carcinoma. This technique also revealed a variant splicing form of TADG-12 that could lead to a truncated protein product. Semi-quantitative PCR showed that TADG-12 is overexpressed in 41 of 55 ovarian cancer specimens relative to normal expression, and the variant form, TADG-12V is found at increased levels in 8 of 22 carcinomas examined. Northern blot revealed three transcripts, the largest of which is approximately 2.4 kb. An ovarian tumor cDNA library was screened, and the entire cDNA of TADG-12 has been identified. This sequence encodes a putative protein of 454 amino acids which includes a potential transmembrane domain, an LDL receptor-like domain, a scavenger receptor cysteine-rich domain, and a serine protease domain. These features imply that TADG-12 will be at the cell surface, and it may be useful as a molecular target for therapy or a diagnostic marker.

PMID: 11068177 [PubMed - indexed for MEDLINE]

Display  Show  Sort by  Send to

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)